

Unimodality of generalized Gaussian coefficients.

Anatol N. Kirillov

*Steklov Mathematical Institute,
Fontanka 27, St.Petersburg, 191011, Russia*

January 1991

Abstract

A combinatorial proof of the unimodality of the generalized q -Gaussian coefficients $\left[\begin{smallmatrix} N \\ \lambda \end{smallmatrix} \right]_q$ based on the explicit formula for Kostka-Foulkes polynomials is given.

1⁰. Let us mention that the proof of the unimodality of the generalized Gaussian coefficients based on theoretic-representation considerations was given by E.B. Dynkin [1] (see also [2], [10], [11]). Recently K.O'Hara [6] gave a constructive proof of the unimodality of the Gaussian coefficient $\left[\begin{smallmatrix} n+k \\ n \end{smallmatrix} \right]_q = s_{(k)}(1, \dots, q^k)$, and D. Zeilberger [12] derived some identity which may be consider as an "algebraization" of O'Hara's construction. By induction this identity immediately implies the unimodality of $\left[\begin{smallmatrix} n+k \\ n \end{smallmatrix} \right]_q$. Using the observation (see Lemma 1) that the generalized Gaussian coefficient $\left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q$ may be identified (up to degree q) with the Kostka-Foulkes polynomial $K_{\tilde{\lambda}, \mu}(q)$ (see Lemma 1), the proof of the unimodality of $\left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q$ is a simple consequence of the exact formula for Kostka-Foulkes polynomials contained in [4]. Furthermore the expression for $K_{\tilde{\lambda}, \mu}(q)$ in the case $\lambda = (k)$ coincides with identity (KOH) from [8]. So we obtain a generalization and a combinatorial proof of (KOH) for arbitrary partition λ .

2⁰. Let us recall some well known facts which will be used later. We base ourselves [9] and [5]. Let $\lambda = (\lambda_1 \geq \lambda_2 \geq \dots)$ be a partition, $|\lambda|$ be the sum of its parts λ_i , $n(\lambda) = \sum_i (i-1)\lambda_i$ and $\left[\begin{smallmatrix} n \\ \lambda \end{smallmatrix} \right]_q$ be the generalized Gaussian coefficient.

Recall that

$$s_\lambda(1, \dots, q^n) = q^{n(\lambda)} \left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q = \prod_{x \in \lambda} \frac{1 - q^{n+c(x)}}{1 - q^{h(x)}}. \quad (1)$$

Here $c(x)$ is the content and $h(x)$ is the hook length corresponding to the box $x \in \lambda$, [5].

Lemma 1 *Let λ be a partition and fix a positive integer n . Consider new partitions $\tilde{\lambda} = (n \cdot |\lambda|, \lambda)$ and $\mu = (|\lambda|^{n+1})$. Then*

$$q^{\frac{n(n-1)}{2} \cdot |\lambda| + n(\lambda)} \left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q = K_{\tilde{\lambda}, \mu}(q). \quad (2)$$

Proof. We use the description of the polynomial $q^{|\lambda|+n(\lambda)} \cdot \left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q$ as a generating function for the standard Young tableaux of the shape λ filled with numbers from the interval $[1, \dots, n]$. Let us denote this set of Young tableaux by $STY(\lambda, \leq n)$. Then

$$q^{|\lambda|+n(\lambda)} \left[\begin{smallmatrix} n \\ \lambda' \end{smallmatrix} \right]_q = \sum_{T \in STY(\lambda, \leq n)} q^{|T|}. \quad (3)$$

Here $|T|$ is the sum of all numbers filling the boxes of T . For any tableau T (or diagram λ) let us denote by $T[k]$ (or $\lambda[k]$) the part of T (or λ) consisting of rows starting from the $(k+1)$ -st one. Given tableau $T \in STY(\lambda, \leq n)$, then consider tableau $\tilde{T} \in STY(\tilde{\lambda}, \mu)$ such that $\tilde{T}[1] = T + \text{supp } \lambda[1]$, and we fill the first row of $\tilde{T}$ with all remaining numbers in increasing order from left to right. Here for any diagram λ we denote by $\text{supp } \lambda$ the plane partition of the shape λ and content $(1^{|\lambda|})$. It is easy to see that

$$c(\tilde{T}) = |T| + \frac{(n+1)(n-2)}{2} \cdot |\lambda|,$$

so we obtain the identity (2). ■

Let us consider an explanatory example. Assume $\lambda = (2, 1)$, $n = 3$. Then $\tilde{\lambda} = (9, 2, 1)$, $\mu = (3^4)$. It is easy to see that $|STY(\lambda, \leq 3)| = 8$.

T	$ T $	$\tilde{T}$	$c(\tilde{T})$
$\begin{smallmatrix} 1 & 1 \\ 2 \end{smallmatrix}$	4	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 3 & 3 & 4 & 4 & 4 \\ 2 & 2 & & & & & & & \\ 3 & & & & & & & & \end{smallmatrix}$	10
$\begin{smallmatrix} 1 & 1 \\ 3 \end{smallmatrix}$	5	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 3 & 3 & 3 & 4 & 4 \\ 2 & 2 & & & & & & & \\ 4 & & & & & & & & \end{smallmatrix}$	11
$\begin{smallmatrix} 1 & 2 \\ 2 \end{smallmatrix}$	5	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 4 & 4 & 4 \\ 2 & 3 & & & & & & & \\ 3 & & & & & & & & \end{smallmatrix}$	11
$\begin{smallmatrix} 1 & 2 \\ 3 \end{smallmatrix}$	6	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 3 & 4 & 4 \\ 2 & 3 & & & & & & & \\ 4 & & & & & & & & \end{smallmatrix}$	12
$\begin{smallmatrix} 1 & 3 \\ 2 \end{smallmatrix}$	6	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 3 & 4 & 4 \\ 2 & 4 & & & & & & & \\ 3 & & & & & & & & \end{smallmatrix}$	12
$\begin{smallmatrix} 1 & 3 \\ 3 \end{smallmatrix}$	7	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 3 & 3 & 3 & 4 \\ 2 & 4 & & & & & & & \\ 4 & & & & & & & & \end{smallmatrix}$	13
$\begin{smallmatrix} 2 & 2 \\ 3 \end{smallmatrix}$	7	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 2 & 3 & 4 & 4 \\ 3 & 3 & & & & & & & \\ 4 & & & & & & & & \end{smallmatrix}$	13
$\begin{smallmatrix} 2 & 3 \\ 3 \end{smallmatrix}$	8	$\begin{smallmatrix} 1 & 1 & 1 & 2 & 2 & 2 & 3 & 3 & 4 \\ 3 & 4 & & & & & & & \\ 4 & & & & & & & & \end{smallmatrix}$	14

Now we would like to use the formula for Kostka-Foulkes polynomials, obtained in [4].

3⁰. First let us recall some definitions from [4]. Given a partition λ and composition μ , a configuration $\{\nu\}$ of the type (λ, μ) is, by definition, a collection of partitions $\nu^{(1)}, \nu^{(2)}, \dots$ such that

- 1) $|\nu^{(k)}| = \sum_{j \geq k+1} \lambda_j$;
- 2) $P_n^{(k)}(\lambda, \mu) := Q_n(\nu^{(k-1)}) - 2Q_n(\nu^{(k)}) + Q_n(\nu^{(k+1)}) \geq 0$ for all $k, n \geq 1$,
where $Q_n(\lambda) := \sum_{j \leq n} \lambda'_j$, and $\nu^{(0)} = \mu$.

Proposition 1 [4] *Let λ be a partition and μ be a composition, then*

$$K_{\lambda, \mu}(q) = \sum_{\{\nu\}} q^{c(\nu)} \prod_{k, n} \left[\begin{matrix} P_n^{(k)}(\lambda, \mu) + m_n(\nu^{(k)}) \\ m_n(\nu^{(k)}) \end{matrix} \right]_q, \quad (4)$$

where the summation in (4) is taken over all configurations of $\{\nu\}$ of the type (λ, μ) , $m_n(\nu^{(k)}) = (\nu^{(k)})'_n - (\nu^{(k)})'_{n+1}$.

From Lemma 1 and Proposition 1 we deduce

Theorem 1 *Let λ be a partition. Then*

$$\left[\begin{matrix} N \\ \lambda' \end{matrix} \right]_q = \sum_{\{\nu\}} q^{c_0(\nu)} \prod_{k, n} \left[\begin{matrix} P_n^{(k)}(\lambda, N) + m_n(\nu^{(k)}) \\ m_n(\nu^{(k)}) \end{matrix} \right]_q, \quad (5)$$

where the summation in (5) is taken over all collections $\{\nu\}$ of partitions $\{\nu\} = \{\nu^{(1)}, \nu^{(2)}, \dots\}$ such that

$$1) |\nu^{(k)}| = \sum_{j \geq k} \lambda_j, \quad k \geq 1, \quad |\nu^{(0)}| = 0,$$

2) $P_n^{(k)}(\nu, N) := n(N+1) \cdot \delta_{k,1} + Q_n(\nu^{(k-1)}) - 2Q_n(\nu^{(k)}) + Q_n(\nu^{(k+1)}) \geq 0$,
for all $k, n \geq 1$. Here

$$c_0(\nu) = n(\nu^{(1)}) - n(\lambda) + \sum_{k, n \geq 1} \left(\alpha_n^{(k)} - \frac{\alpha_n^{(k+1)}}{2} \right), \quad \alpha_n^{(k)} := (\nu^{(k)})'_n \quad (6)$$

and by definition $\left(\begin{matrix} \alpha \\ 2 \end{matrix} \right) := \frac{\alpha(\alpha-1)}{2}$ for any $\alpha \in \mathbf{R}$.

The identity (5) may be considered as a generalization of the (KOH) - identity (see [8]) for arbitrary partition λ .

Corollary 1 *The generalized q -Gaussian coefficient $\left[\begin{matrix} n \\ \lambda' \end{matrix} \right]_q$ is a symmetric and unimodal polynomial of degree $(N-1)|\lambda| - n(\lambda)$.*

Proof. First, it is well known (e.g. [10],[11]) that the product of symmetric and unimodal polynomials is again symmetric and unimodal. Secondly, we use a well known fact (e.g.[10]), that the ordinary q -Gaussian coefficient $\left[\begin{matrix} m+n \\ n \end{matrix} \right]_q$ is a symmetric and unimodal polynomial of degree mn . So in order to prove Corollary 1, it is sufficient to show that the sum

$$2c_0(\nu) + \sum_{k,n} m_n(\nu^{(k)}) P_n^{(k)}(\nu, N) \quad (7)$$

is the same for all collections of partitions $\{\nu\}$ which satisfy the conditions 1) and 2) of the Theorem 1. In order to compute the sum (7), we use the following result (see [4]):

Lemma 2 Assume $\{\nu\}$ to be a configuration of the type (λ, ν) . Then

$$\sum_{k,n} m_n(\nu^{(k)}) P_n^{(k)}(\nu, \mu) = 2n(\mu) - 2c(\nu) - \sum_{n \geq 1} \mu'_n \cdot \alpha_n^{(1)}. \quad (8)$$

Using Lemma 2, it is easy to see that the sum (7) is equal to $(N-1)|\lambda| - n(\lambda)$. This concludes the proof. $\blacksquare$

Note that in the proof of Corollary 1 we use symmetry and unimodality of the ordinary q -Gaussian coefficient $\left[\begin{matrix} m+n \\ n \end{matrix} \right]_q$. However, we may prove

the unimodality of $\left[\begin{matrix} m+n \\ n \end{matrix} \right]_q$ by induction using the identity (5) for the case $\lambda = (1^n)$, $N = m$.

Remark 1. The unimodality of generalized q -Gaussian coefficients was also proved in the recent preprint [7]. The proof in [7] uses the result from [4]. However [7] does not contain the identity (5).

Remark 2. The proof of the identity (4) given in [4] is based on the construction and properties of the bijection (see [4])

$$STY(\lambda, \mu) \rightleftharpoons QM(\lambda, \mu).$$

It is an interesting task to obtain an analytical proof of (5). In the case $q = 1$ such a proof was obtained in [3].

Acknowledgements. The final version of this paper was written during the author's stay at RIMS. The author expresses his deep gratitude to RIMS for its hospitality.

References

- [1] Dynkin E.B., Some properties of the weight system of a linear representation of a semisimple Lie group (in Russian). Dokl. Akad. Nauk USSR, 1950, 71, 221-224.
- [2] Hughes J.W., Lie algebraic proofs of some theorems on partitions. *In* Number Theory and Algebra, Ed. H. Zassenhaus, Academic Press, NY, 1977, 135-155.
- [3] Kirillov A.N., Completeness of the Bethe vectors for generalized Heisenberg magnet. Zap. Nauch. Sem. LOMI (in Russian), 1984, 134, 169-189.
- [4] Kirillov A.N., On the Kostka-Green-Foulkes polynomials and Clebsch-Gordan numbers. Journ. Geom. and Phys., 1988, 5, 365-389.
- [5] Macdonald I.G., Symmetric Functions and Hall Polynomials. Oxford University Press, 1979.
- [6] O'Hara K., Unimodality of Gaussian coefficients: a constructive proof. Jour. Comb. Theory A, 1990, 53, 29-52.
- [7] Goodman F., O'Hara K., Stanton D., A unimodality identity for a Schur function. Preprint 1990/1991.
- [8] Stanton D., Zeilberger D., The Odlyzko conjecture and O'Hara's unimodality proof. Bull. Amer. Math. Soc., 1989, 107, 39-42.
- [9] Stanley R., Theory and application of plane partitions I,II. Studies Appl. Math., 1971, 50, 167-188, 259-279.
- [10] Stanley R., Unimodal sequences arising from Lie algebras. *In* Young Day Proceedings, Eds. J.V.Narayana, R.M.Mathsen, and J.G.Williams, Dekker, New York/Basel, 1980, 127-136.
- [11] Stanley R., Log-concave and unimodal sequences in algebra, combinatorics, and geometry. Annals of the New York Academy of Sciences, 1989, 576, 500-535.
- [12] Zeilberger D., Kathy O'Hara's constructive proof of the unimodality of the Gaussian polynomials. Amer. Math. Monthly, 1989, 96, 590-602.